Heraklion (Crete), August 22, 2010



Session 6 - Equine milk for human health

Mechanical milking of equines, the european proposal

Simoni A.1, Drogoul C.2, Salimei E.3

1 Di SAVA, Università degli Studi del Molise, Campobasso (IT)

2 LEGTA Semur Chatillon, Chatillon sur Seine (F)

3 Di. STAAM, Università degli Studi del Molise, Campobasso (IT)






Portable milking machine for mares

(Le Du, 1986 photo Faubladier, 2005)








(photo Salimei 1996)



Milk ejection in mare



Le Du, 1986

MILK FRACTION	FAT CONTENT, %
Cisternal milk	0.15
Alveolar milk	0.5 - 1.1
Stripping	1.1 - 1.4
Residual milk (~ 30% total yield)	3.5 – 7.2


Characteristics of portable milking machine (Le Du, 1986)

Pulsation rate	120 cycles/min
Pulsation ratio	50%
Vacuum level	42 kPa (37 ÷50 kPa)
Cluster (unit)	Sheep or goat type
Bucket	Pour milk gently in order to avoid physical stress

Dairy mare – machine milking management (Bayle-Labouré, 2007)


Milking frequency	Three times per day
Beginning of milking period	2 – 3 months after foaling
Milking period	From june to december (6 -7 months)
Milking routine	3 hr physical separation from foalmassage udder and manual stimulationcleaning teat
Foal management	Foals nurse at night and during week-ends
Milk yield	2 ÷ 2.5 1/ milking (heavy breed) about 1000 1/mare·season
Facilities	Milking parlour close to milk room

Milking stable for mare



From Bayle-Labouré, 2007

Mare's milk production



Doreau and Martin Rosset, 2002



Milking facility in a dairy donkey farm



Project: Simoni, 2004; prototype Borghi



Dairy donkey farm: a proposal for marginal areas





Simoni, 2010



Photo: www.donkeywelfare.com

Andrea, Carole and Elisabetta thank you for your attention