



Estimation of NDF degradation parameters in practice



REDNEX

Martin Riis Weisbjerg

**Faculty of Agricultural Sciences,
Aarhus University, PO Box 50,
Tjele, Denmark**



***Innovative and practical management
approaches to **reduce nitrogen**
excretion by ruminants***

Some explanations:

Ruminants


Estimation of **NDF degradation parameters** in practice

NDF – Neutral detergent fibre

Degradation vs. digestion

Degradation parameters

- Potential degradability
- Rate of degradation, k_d



Content:

**New feed evaluation systems for ruminants –
challenge to feed evaluation**

Importance of NDF in feed rations

Physical structure

Feed intake

Energy supply

Estimation of NDF degradability

Research methods

Practical methods

Conclusion

New feed/ration evaluation systems for ruminants

Understanding of the ruminal ecosystem has shown the shortcomings of the classic additive feed evaluation systems

Future feed/ration evaluation tools should improve simulation/prediction/monitoring of nutrients available, to predict and optimise production

Examples of ration formulation systems:

- CNCPS (Cornell net carbohydrate and protein system)**
- NorFor (Nordic countries)**

The challenge - feed evaluation



Feed evaluation will still be based on individual feeds

The challenge → provide data for potential digestibility and rate of digestion for main nutrients

Need for tabulated values as default values


Challenge is to develop analytical tools for estimation of degradation parameters on samples from practical agriculture

This presentation will focus on NDF


Based on:

- Experience from ‘NorFor Feed table working group’**
- Results and plans - RedNex WP1 project - Tools for feedstuff evaluation to predict protein supply in dairy cows**

What is NDF? The residue not solubilised after boiling with a neutral detergent solution



Degradation parameters




$$\text{Degradation} = \text{dNDF}(1 - e^{-k_d t})$$

NDF digestibility main factor for milk production

- Classical feed evaluation systems failed

Observed - expected yield



Physical structure:

NDF gives the physical structure in feed

Chewing, rumination

Rumen motility

Rumen environment

Rumen mat – rumen stratification

But is all NDF equal?

Roughage vs. concentrate

Particle size (Physical effective NDF >1.18 mm)

Digestibility – indigestibility

Associative effects of ration, on digestibility etc.

Feed intake:

NDF gives the bulk of the ration – therefore ration NDF concentration and digestibility are the main factors determining *ad libitum* feed intake in a physical regulated ruminant

Energy supply:

NDF is the largest individual nutrient fraction in most feed rations to ruminants (30% of DM in Danish rations)


Organic matter (OM) digestibility is determined by NDF

Because:

- Cell content (NDS, neutral detergent solubles) true digestibility ~ 100**
- Thereby variation in OM digestibility is due to variation in NDF concentration and digestibility**

Therefore, rate and extent of NDF degradation is of outmost importance for the energy supply to the ruminant – increased by the effect on feed intake

Rumen digestibility – competition between digestion and passage



$$\text{Digestibility} = k_d / (k_d + k_p)$$


Methods – digestibility and degradation




In vivo – feed - faeces difference


In vitro - solubility after in vitro treatment with rumen fluid or commercial enzymes

In situ – degradation after feed has been incubated in the rumen in nylon/dacron bags with pores





Examples of NDF degradation profiles – fresh and ensiled grass and grass/clover




Estimation of NDF degradability

Research methods

Main research methods

Rate of degradation (k_d)


- In situ degradability
- In vitro degradability
- In vitro gas production





Potential degradability (dNDF, iNDF)

- In situ degradability
- In vitro degradability

Effect of methods (in situ, in vitro, pH) on NDF degradation profiles



**In situ vs. gas
production – Test
against in vivo data**
- Both methods
predict in vivo
digestibility well



Conclusions k_d

All methods have in build problems

Difficult to say some are better, some worse

Very few tests on in vivo data!!!!!!

Potential degradability (dNDF, iNDF)

- In situ degradability
- In vitro degradability
- Estimation based on residue after long time incubation
- Main problems:
 - Particle loss in situ
 - Maintaining fermentation in vitro

Estimation of NDF degradability

Practical feed evaluation



September 10, 2007

NorFor in sacco standard

At the seminar "Laboratory methods to predict in situ degradation profiles" held in Uppsala November 17, 2004, methodological aspects on in sacco determination of rumen degradability were discussed by invited scientists and the NorFor Feed Table Group. One important goal with the seminar was to standardize the in sacco procedure as much as possible to minimize between-laboratory variation. Critical parts of the method were listed at the seminar and completed by literature review of other published standards (Madsen & Hvelplund, 1994; Madsen et al., 1995; VanZant et al., 1998; IAEA, 2000; NRC, 2001) and papers on methodological details (Lindberg, 1985; De Boer et al., 1987; Cherney et al., 1990; Varvikko and Vanhatalo, 1990; Uden, 1992; Madsen and Hvelplund, 1994; Wilkerson et al., 1995; Coblenz et al., 1997; Huntington and Givens, 1997a; Huntington and Givens, 1997b; Huntington and Givens, 1997c). Preliminary proposals for the standard have been modified by the NorFor Feed Table Group after consulting scientists that attended the Uppsala seminar. The standard presented in Table 1 is the final agreement of the NorFor Feed Table Group.

The Feed Table Group in NorFor:

Torsten Eriksson, Swedish University of Agricultural Sciences

Erica Lindberg, Swedish Dairy Association

Odd-Magne Harstad, Norwegian University of Life Sciences (UMB)

Lars Bævre, TINE, Norway

Bragi Lindal Olafsson, Agricultural Research Institute, Iceland

Martin Weisbjerg, University of Aarhus

Rudolf Thøgersen, Danish Agricultural Advisory Service, National Centre, Danish Cattle Federation

Table 1. NorFor In sacco standard

Item	NorFor standard
Animal	
Type	Dry cow, dairy breed, representative animal
Feeding level	Maintenance
Diet	(Hay+straw):concentrate 67:33. CP content of ration DM >12%. The concentrate should contain a minimum of 3 sources of protein
Meals	Daily ration should be divided in 2 or more meals of equal size
Minimum adaptation period to diet	14 days but if the animal has been on pasture or otherwise been fed on a diet and level totally different from the standard, minimum adaptation period is 21 days
Replication	
Number of animals	3 cows except for INDF determination where 2 cows is sufficient
Bags per animal	Not specified
Number of days when sample is replicated	1 (=days are not replicated)
Sample preparation	
Drying	Freeze-drying preferable but oven drying at 45° C also allowed. For NDF determination, a drying temperature of 60° C is allowed
Grinding	Screen aperture 1.5 mm. Cutter mill preferable but hammer mill allowed during NorFor's introduction phase
Sample size	1.0 – 2.0 grams dried sample. See "Sample size to surface area" below

NorFor

In situ standard

Feed table

<http://www.norfor.info/>

Methods to be used in practice

Rate of degradation (k_d)

- In situ degradability
- In vitro degradability
- In vitro gas production
- NIRs
- Multiple regressions on chem. and dig. measures
- Backwards calculation

Potential degradability (dNDF, iNDF)

- In situ degradability
- In vitro degradability
- NIRs
- Multiple or simple regression (chem., dig.)

Rate of degradation (k_d)



NIRs


Limited success predicting rate of degradation for both NDF and other nutrients

More efficient in predicting solubilities and potential digestibilities

Rate of degradation (k_d)

Regressions on chemical or digestibility measurements

On grass – grass/clover, possible to explain 86% of variation in k_d by in vitro enz. NDF digestibility



Koukolová et al., J. Anim Feed Sci. 2004

Also high for barley and wheat whole crops, 0.81 and 0.77, respectively, to in vitro enz. OM digestibility

Weisbjerg, M.R., Mikkelsen, M., Bossen, D. & Lund, P. J. Dairy Sci. 2003

Rate of degradation (k_d)



'Backwards' calculation

Information needed

OM digestibility

Ash concentration

NDF concentration

iNDF concentration

All except iNDF classical feed analysis

Idea:

NDS digestibility estimated using Lucas principle

NDF digestibility calculated by difference

Kd NDF 'backwards' calculated assuming 2 pool rumen model

Potential degradability (dNDF, iNDF)




iNDF in feeds in practice

- Large variations in values
 - Between feedstuff groups
 - Within group
 - Within feedstuff type (maturity, processing etc.)

Potential degradability (dNDF, iNDF)



iNDF vs. ADL



Potential degradability (dNDF, iNDF)



NIRs calibration

**In the Nordic countries we pt. use calibrations
calibrated directly on in situ iNDF**

Potential degradability (dNDF, iNDF)



**Lack of good, reliable and cheap lab methods,
for NIRs calibration**

**iNDF= 2.4 x ADL (CNCPS ratio) only fits for maize
silage, barley whole crop, lucerne, wheat**

**ADL content and/or IVOMD acceptable predictors of
iNDF within feedstuff group**

Important research area in coming years

Conclusions



New feed/ration evaluation systems require cheap/efficient methods for estimation of NDF degradability

- **Rate of degradation**
- **Potential degradability**

Research methods available, however quality and in vivo documentation problematic!

Practical methods:

Rate of degradation

NIRs problematic

Simple regressions useful within feedstuff type

The backwards calculation might be the future

Potential degradability

NIRs or similar 'cheap' methods the future

However, reliable laboratory methods needed for NIRs calibration

This presentation has been carried out with financial support from the Commission of the European Communities, FP7, KBB-2007-1.



**It does not necessarily reflect its view
and in no way anticipates the
Commission's future policy in this area.**




***Innovative and practical management
approaches to **reduce nitrogen**
excretion by ruminants***



kdNDF in Forage

The backward calculation method (Weisbjerg et al 2004, 2007)



$$ED = \left(\frac{kd}{(kd + kr)} \cdot \left(1 + \frac{kr}{(kd + kp)} \right) \right) \cdot Pd$$

Allen & Mertens, 1988

$$kdNDF = 1,21 \cdot 100 \cdot \left(-0,041667 + \frac{\sqrt{0,006944 + \frac{0,0066667 \cdot D}{1 - D}}}{2} \right) - 1,24$$

Huhtanen et al, 2006

$$NDS = 1000 - Ash - NDF$$

$$NDSdig = (101,3 - (902/NDS/10))/100$$

$$uOM = (1000 - Ash) \cdot (1 - \text{OMD}/100)$$

$$uNDS = NDS \cdot (1 - NDSFK)$$

$$uNDF = uOM - uNDS$$

$$NDFdig = (NDF - uNDF)/NDF$$

$$pdNDF = NDF - iNDF$$

$$D = NDFdig/(pdNDF/1000)$$

**OMD
estimated
from sheep
fed at
maintenance**