

Comparison Difference Levels of Phytase Enzyme in Diet of Japanese Quail (*Coturnix japonica*) and Some Blood Parameters

Nasrollah Vali¹

Department of Animal Sciences, faculty of Agriculture
Islamic Azad University Shahrekord Branch, Iran
¹nasrollah.vali@gmail.com

Abstract

A total of 210, 3-d-old Japanese quail chicks, (*Coturnix japonica*) allocated to 21 cages, each cage containing 10 chicks, that were received seven diets with three replication of each diet. Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments 1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls). At the end of each experimental period (each week to 45 days of ages), birds were weighed and feed consumption was recorded for feed efficiency computation. At the age of 45 days, four chicks were randomly selected from each treatment blood samples for subsequent determination of minerals (Ca, P and Mg) and alkaline phosphates (ALP) in serum. Data were subjected to analysis of variance using the general linear models (GLM) procedures of SAS software. Data showed that feed efficiency was not affected at periods of 9-16 and 37- 45 days of ages by addition of phytase, but in other periods were significantly different ($P < 0.05$). Liver weight in treatment 5 (4.68 ± 0.23 gr) at 45 days of age was more than other groups that had significant difference ($P < 0.05$). Quails of treatment 4 had Calcium (Ca) and Phosphorus (P) 10.19 ± 0.80 , 7.11 ± 0.46 mg/dl respectively that were more than other groups. The amount of Magnesium on treatment 1 (3.27 ± 0.15 gr), positive control (3.27 ± 0.15 gr) and negative control (3.56 ± 0.15 gr) which were not significantly different ($P > 0.05$), whereas treatment 1 and control groups had significant differences with other treatments ($P < 0.05$). The Alkaline phosphates of treatment 5 (366.83 ± 38.07 IU/L) in blood serum at 45 days of age represents the most increase among other treatments, which had significant differences with treatment 1 (124.99 ± 50.80 IU/L) and control groups, although weren't significant difference with the treatments 2 (260.26 ± 50.01 IU/L) and treatments 3 (288.37 ± 40.64 IU/L).

Key words: Blood Factors, *Coturnix japonica*, Phytase Enzyme, quail

Introduction

The primary constituents of diets for poultry are plant-based ingredients which come primarily from the seeds of plants. Most of the stored phosphorus in plants is found in seeds mainly as a component of a molecule called phytin. Phytin-phosphorus is poorly available to poultry, and this availability varies both within and among ingredients. The enzyme phytase releases phosphate groups from phytin potentially making this released phosphorus available to the animal. Phytase is the only recognized enzyme that can initiate the release of phosphate from phytin (International Union of Biochemistry, 1979). Nelson et al., (1968) demonstrated that the addition of the enzyme phytase to grains and feeds was an effective way to increase phosphorus (P) availability to poultry.

Numerous reports have shown that phytate phosphorus present in cereal grains could be made available to both poultry and swine by treating the grains or supplementing the diets with the enzyme phytase, reducing the need to supplement these diets with inorganic P and decreasing P excretion (Jongbloed and Kemme, 1990; Simons *et al.*, 1990; Zhu *et al.*, 1990; Ketaren *et al.*, 1993).

P is an essential mineral for laying hens in formation of egg shell and metabolism (Frost and Roland, 1991; Summers, 1995; Usayran and Balnave, 1995). Only 20-50% of plant P is available to broilers, and the rest of P is in the form of phytate (myo-inositol hexaphosphate), which is poorly used by broilers (Ravindran *et al.*, 1995). Ravindran *et al.*, (1998) and Sebatian *et al.*, (1998) reported that poultry can not produce enough amounts of endogenous phytase to hydrolyze and release P from phytate. To meet dietary P requirement of laying hens, inorganic P (Dicalcium phosphate) or exogenous phytase enzymes are commonly added to commercial corn-soy layer diets. However, Dicalcium phosphate supplementation is not only expensive but also leads to environmental problems by over-supplementation. Excess P from the feces of hens is easy to access ground water, rivers, lakes, and oceans, and can lead to mortality of aquatic animals by stimulating algae growth.

Considerable research has demonstrated that phytase supplementation (from 100 to 300 phytase unit (FTU)/kg of feed) to diets containing 0.1% dietary nonphytate phosphorus (NPP) has positive effects on egg production, egg mass, egg weight, egg specific gravity, bone ash, and eggshell quality by improving P utilization (Boiling *et al.*, 2000a, b; Jalal and Scheideler, 2001; Roland *et al.*, 2003; Keshavarz, 2003). Phytase supplementation decreased P excretion in the manure and reduced the potential environmental problems (Jalal and Scheideler, 2001; Biehl and Baker, 1997).

Through consideration authentic resources the phytase enzyme was narrow in diets of quail so, the target of this study is comparison difference levels of Phytase enzyme in diet of Japanese quail (*Coturnix japonica*) for investigation of feed consumption and some blood factors.

Materials and methods

A total of 210, 3-d-old Japanese quail chicks, (*Coturnix japonica*) were housed in pen with the newspaper litter until 12 days of age. Then birds were moved into one-tier cage (60×50×40 cm) in an environmentally controlled room, with 23-h constant overhead fluorescent lighting. The temperature was decreased gradually from 35°C to 20°C on the weekly bases. The chicks were allocated to 21 cages, each cage containing 10 chicks, that were received seven diets with three replication of each diet. The experimental diets are presented in Table 1 and were formulated to meet the requirements of quail as established by the NRC (1994). Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls). One FTU is defined as the amount of enzyme that liberates micromole inorganic phosphorus per minute from 0.0051 mol/L sodium phytate at 37° C and at pH = 5.5 Hall *et al.*, (2003). Small amounts of the basal diet were first mixed with the respective amounts of phytase as a small batch and then with a larger amount of the basal diets until the total amounts of the respective diets were homogeneously mixed. Diets in mash form and water were provided ad libitum from 3 to 45 days of ages. At the end of each experimental period (each week to 45 days of ages), birds were weighed and feed consumption was recorded for feed efficiency computation.

At the age of 45 days, four quails were randomly selected from each treatment and weighed, and were killed, bled and plucked for liver weight and blood samples for subsequent determination of minerals (Ca, P and Mg) and alkaline phosphates (ALP) in serum. A total 84 blood samples (8-10 cc) collected for to assigning Ca, p, Mg elements by photometric system and alkaline phosphates enzyme (ALP) by photometric system by method Deutsche Gesellschaft für klinische Chemie (1972).

Data were subjected to analysis of variance using the general linear models procedures of SAS software (SAS Institute, 1998). Statements of statistical significance are based on a probability of $P < 0.05$.

Table 1: Ingredients and nutrient composition of the experimental diets

Items	Phytase Levels (FTU/Kg) [†]						
	0	0					
Ingredients (%)	(positive control)	(negative control)	150	300	600	1200	2400
Corn (8.5% CP)	49	49	49	49	49	49	49
Soybean meal(44% CP)	44.91	44.91	44.91	44.91	44.91	44.91	44.91
Vegetable oil (8.8 kcal/g)	2.865	2.865	2.865	2.865	2.865	2.865	2.865
Limestone	1.304	1.304	1.304	1.304	1.304	1.304	1.304
Salt (NaCl)	0.20	0.20	0.20	0.20	0.20	0.20	0.20
Dicalcium phosphate	0.747	0.212	0.212	0.212	0.212	0.212	0.212
DL-Methionine	0.136	0.136	0.136	0.136	0.136	0.136	0.136
Mix vitamins [‡]	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Trace mineral [§]	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Sand	0.337	0.873	0.870	0.867	0.861	0.849	0.825
Phytase (5000 FTU/ g)	0	0	0.003	0.006	0.012	0.024	0.048
Total	100	100	100	100	100	100	100
Nutrient (Calculated)							
Metabolizable energy (Kcal/Kg)	2900	2900	2900	2900	2900	2900	2900
Crude protein (%)	24	24	24	24	24	24	24
Methionine + cystine (%)	0.884	0.884	0.884	0.884	0.884	0.884	0.884
Lysine (%)	1.33	1.33	1.33	1.33	1.33	1.33	1.33
Calcium (%)	0.800	0.682	0.682	0.682	0.682	0.682	0.682
Total phosphorus (%)	0.569	0.469	0.469	0.469	0.469	0.469	0.469
Available phosphorus (%)	0.300	0.200	0.200	0.200	0.200	0.200	0.200

[†] Danisco Animal Nutrition, Carol Stream, IL, FTU/kg feed = phytase units/kg feed. One phytase unit (FTU) is defined as the amount of enzyme that liberates micromole inorganic phosphorus per minute from 0.0051 mol/L sodium phytate at 37° C and at pH = 5.5.

[‡] Provided per kilogram of diet: vitamin A (as retinyl acetate), 8,000 IU; cholecalciferol, 2,200 ICU; vitamin E (as DL- α -tocopheryl acetate), 8 IU; vitamin B12, 2800.02 mg; riboflavin, 5.5 mg; D-calcium pantothenic acid, 13 mg; niacin, 36 mg; choline, 500 mg; folic acid, 0.5 mg; vitamin B1 (thiamin mononitrate), 1 mg; pyridoxine, 2.2 mg; d-biotin, 0.05 mg; vitamin K (menadione sodium bisulfate complex), 2 mg.

[§] Trace mineral mix provided the following (per kg of diet): manganese (MnSO₄·H₂O), 60 mg; iron (FeSO₄·7H₂O), 30 mg; zinc (ZnO), 50 mg; copper (CuSO₄·5H₂O), 5 mg; iodine (ethylene diamine dihydroiodide), 1.5 mg.

Results and discussion

The effects of phytase enzyme on growth performance are summarized in Table 2. The main effects data indicated that the body weights at 23 and 45 days of ages weren't significantly different, and at the other ages (16, 30 and 37 days of age) were significantly different, but in all stages positive control with other experiment treatments weren't significantly different ($P < 0.05$). Body weight at 45 days of age in treatment of 5 was 173.77 ± 13.40 gr that showed more than other treatments, and treatment of 3 in the same age with 156.00 ± 13.40 gr was the lowest among other treatments (Table 2). Data also showed that feed efficiency (Table 3) was not affected at periods of 9-16 (FE1) and 37- 45 (FE5) days of ages by addition of phytase, as a result of simultaneous showed in weight gain, but in other periods were significantly different ($P < 0.05$). In this study, feed efficiency weren't different between the positive and negative control groups (Table 3) in any periods ($P < 0.05$).

These results were similar to those obtained by Perney *et al.*, (1993), Sebastian *et al.*, (1996), Ahmad *et al.*, (2000) and Viveros *et al.*, (2002) those have reported on broiler. Although, Rama-Rao *et al.*, (1999) reported improved feed efficiency with supplemental phytase. These contrasting results may be due to a number of factors including phytase source, ingredients (type, source, phytate content), and dietary characteristics (processing, Vitamin D3 level, Ca:P ratio) (Ravindran *et al.*, 1995a).

Table 2: Least square means for effect treatments and sex with standard error for Body weight average (gr) at age of 16(BW1), 23(BW2), 30(BW3), 37(BW4) and 45(BW5) days of ages.

Treatment ¹	BW1	BW2	BW3	BW4	BW5
1	61.66 ± 3.18^a	83.77 ± 5.00	101.50 ± 6.73^{ab}	133.71 ± 8.75^{ab}	171.86 ± 13.40
2	50.46 ± 3.18^b	74.54 ± 5.00	94.47 ± 6.73^b	122.41 ± 8.75^{ab}	163.91 ± 13.40
3	52.07 ± 3.18^{ab}	70.04 ± 5.00	85.93 ± 6.73^b	115.07 ± 8.75^b	156.00 ± 13.40
4	57.04 ± 3.18^{ab}	77.98 ± 5.00	91.47 ± 6.73^b	112.04 ± 8.75^b	161.01 ± 13.40
5	50.28 ± 3.18^b	73.52 ± 5.00	100.75 ± 6.73^{ab}	135.29 ± 8.75^{ab}	173.77 ± 13.40
Positive control	60.69 ± 3.18^{ab}	83.97 ± 5.00	116.94 ± 6.73^a	146.61 ± 8.75^a	169.15 ± 13.40
negative control	54.29 ± 3.18^{ab}	72.24 ± 5.00	81.35 ± 6.73^b	110.19 ± 8.75^b	165.92 ± 13.40

^{ab} Values with in a column with no common superscript differ significantly ($P < 0.05$) and columns without superscript were not significantly different.

¹ Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments 1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls).

Table 3: Least square means for effect treatments with standard error for feed efficiency(gr/gr) at periods of 9-16 (FE1), 16-23 (FE2), 23- 30 (FE3), 30 37 (FE4) and 37- 45 (FE5) days of ages.

Treatment ¹	FE1	FE2	FE3	FE4	FE5
1	0.96 ± 0.05	1.12 ± 0.04 ^{ab}	1.16 ± 0.05 ^a	1.26 ± 0.03 ^a	1.16 ± 0.07
2	0.95 ± 0.05	1.06 ± 0.04 ^{ab}	1.16 ± 0.05 ^a	1.16 ± 0.03 ^{ab}	1.06 ± 0.07
3	0.98 ± 0.05	1.18 ± 0.04 ^a	1.07 ± 0.05 ^{ab}	1.17 ± 0.03 ^{ab}	1.10 ± 0.07
4	0.89 ± 0.05	1.10 ± 0.04 ^{ab}	1.11 ± 0.05 ^{ab}	1.23 ± 0.03 ^a	1.14 ± 0.07
5	0.87 ± 0.05	1.03 ± 0.04 ^b	1.12 ± 0.05 ^{ab}	1.11 ± 0.03 ^b	1.12 ± 0.07
Positive control	0.85 ± 0.05	1.14 ± 0.04 ^{ab}	0.98 ± 0.05 ^b	1.08 ± 0.03 ^b	1.03 ± 0.07
negative control	0.90 ± 0.05	1.07 ± 0.04 ^{ab}	1.08 ± 0.05 ^a	1.17 ± 0.03 ^{ab}	1.14 ± 0.07

^{ab} Values with in a column with no common superscript differ significantly ($P < 0.05$) and columns without superscript were not significantly different.

¹ Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls).

The effect of these treatments on liver weight (LW), Calcium (Ca), Phosphorus (P), Magnesium (Mg) and alkaline phosphates (ALP) blood serum at 45 days of age are presented in Table 4. Liver weight in treatment 5 (4.68 ± 0.23 gr) at 45 days of age was more than other groups that had significant difference ($P < 0.05$). Quails of treatment 4 had Calcium (Ca) and Phosphorus (P) 10.19 ± 0.80 , 7.11 ± 0.46 mg/dl respectively that were more than other groups. The main effects of data indicated that the increase in phytase content in the diet were not significantly different ($P > 0.05$) for Ca retention at 45 days of age, but for P retention were significantly different ($P < 0.05$) Diagram 1, 2 and Table 4. The amount of Magnesium on treatment 1 (3.27 ± 0.15 gr), positive control (3.27 ± 0.15 gr) and negative control (3.56 ± 0.15 gr) which were not significantly different ($P > 0.05$), whereas treatment1 and control groups had significant differences with other treatments ($P < 0.05$) Diagram 3 and Table 4. The Alkaline phosphates of treatment 5 (366.83 ± 38.07 IU/L) in blood serum at 45 days of age represents the most increase among other treatments, which had significant differences with treatment 1 (124.99 ± 50.80 IU/L) and control groups, although weren't significant difference with the treatments 2 (260.26 ± 50.01 IU/L) and treatments 3 (288.37 ± 40.64 IU/L) Diagram 4 and Table 4. These results were similar to those obtained by Sarcicek and Garipoglu (2005) in laying quails, but more than those obtained by Denek *etal* (2007) in Japanese quails. The decrease in serum alkaline phosphates associated with the diets supplemented with phytase might reflect the down regulation of this enzyme resulting from the increased availability of phosphorus (Huff *etal*.1998).

Phytates (Phytic acid, mineral complexes), as well as very limited providing P availability to the animal in the absence of phytase, can also have other effects which can reduce nutrient digestibility and, consequently, animal performance (i.e. phytates in feed act as anti-nutrients). At neutral or basic pH, phytic acid can form complexes with divalent and trivalent

cations. These minerals and trace elements attach themselves easily onto the phosphate molecule of phytic acid producing insoluble salts. Once these compounds are produced these cations are unavailable for absorption in the gut and, like phosphorus, these valuable nutrients are then also lost to the animal. If these insoluble salts react with fatty acids in the gut, insoluble metallic soaps are produced. It has been suggested that these soaps may lower fat digestibility and consequently may have an effect on energy availability. Phytic acid can also form complexes with proteins and starch. These can occur in the plant (native complexes) or during digestion in the gastrointestinal tract. In the highly acidic stomach region, amino acids are bound directly to phytate phosphorus creating an insoluble complex. In the less acidic intestine, multiple bonded mineral cations act as a bridge between phytate and protein. These complex compounds are more resistant to enzymatic digestion than their unbound counterparts. Moreover, phytates can also potentially decrease protein and starch digestibility by inhibiting, or binding to, digestive enzymes produced by the animal (e.g. pepsin and trypsin for protein degradation and alpha-amylase for starch degradation). The ability of phytase to improve P availability by hydrolyzing phytate-bound P in poultry diets is well documented (Broz *et al.*, 1994; Coelho and Kornegay, 1996; Kornegay *et al.*, 1996; Qian *et al.*, 1996, 1997). Increased utilization of P from phytate can therefore reduce supplementation of diets with inorganic P sources while maintaining normal growth of the bird.

Table 4: Least square means with standard error for Body weight average (BW), liver weight (Lw), Calcium (Ca), Phosphorus (P), Magnesium (Mg) and Alkaline phosphates (ALP) at 45 days of age

Treatments ¹	LW(gr)*	Ca (mg/dl)**	P (mg/dl)	Mg (mg/dl)	ALP (IU/L)***
1	3.58 ± 0.21 ^b	8.20 ± 0.73	4.11 ± 0.46 ^c	3.27 ± 0.15 ^{ab}	124.99 ± 50.80 ^{cd}
2	3.69 ± 0.23 ^b	8.37 ± 0.75	6.32 ± 0.48 ^{ab}	2.40 ± 0.15 ^d	260.26 ± 50.01 ^{abc}
3	3.61 ± 0.23 ^b	9.60 ± 0.73	6.65 ± 0.46 ^{ab}	2.55 ± 0.15 ^{cd}	288.37 ± 40.64 ^{ab}
4	3.49 ± 0.22 ^b	10.19 ± 0.80	7.11 ± 0.46 ^a	2.95 ± 0.16 ^{bc}	200.31 ± 44.34 ^{bcd}
5	4.68 ± 0.23 ^a	10.09 ± 0.79	6.71 ± 0.46 ^{ab}	2.72 ± 0.15 ^{cd}	366.83 ± 38.07 ^a
Positive control	3.90 ± 0.21 ^b	10.08 ± 0.75	3.96 ± 0.46 ^c	3.27 ± 0.15 ^{ab}	131.78 ± 39.84 ^d
Negative control	3.89 ± 0.24 ^b	9.69 ± 0.72	5.86 ± 0.46 ^b	3.56 ± 0.15 ^a	113.36 ± 54.59 ^d

^{abcd} Values with in a column with no common superscript differ significantly ($P < 0.05$) and column without superscript was not significantly different.

¹ Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments 1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls). One FTU is defined as the amount of enzyme that liberates micromole inorganic phosphorus per minute from 0.0051 mol/L sodium phytate at 37°C and at pH = 5.5

* (gr) gram

** (mg/dl) mille gram per deciliter

*** (IU/L) international unit per liter




Diagram 1: Calcium of blood serum (mg/dl) quails at 45 days of age in different treatments.

¹ Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls).




Diagram 2: Phosphorus of blood serum (mg/dl) quails at 45 days of age in different treatments.

¹ Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls).




Diagram 3: Magnesium of blood serum (mg/dl) quails at 45 days of age in different treatments.

¹ Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls).




Diagram 4: Alkaline phosphates of blood serum (IU/L) quails at 45 days of age in different treatments.

¹ Experimental diets were arranged with five levels 150, 300, 600, 1200 and 2400 phytase enzyme unit (FTU/Kg) with commercial name (Natuphos 500) as treatments1, 2, 3, 4 and 5 respectively, and two control groups (positive and negative controls).

Reference:

- Ahmad, T., S. Rasool, M. Sarwar, A. Haq, and Zia-ul Hasan, 2000. Effect of microbial phytase produced from a fungus *Aspergillus niger* on bioavailability of phosphorus and calcium in broiler chickens. *Anim. feed Sci. Techn.*, 83:103–114.
- Biehl, R. R., and D. H. Baker, 1997. Microbial phytase improves amino acid utilization in young chicks fed diets based on soybean meal but not diets based on peanut meal. *Poul. sci.*, 76:355–360.
- Boling, S. D., M. W. Douglas, R. B. Shirley, C. M. Parsons, and K. W. Keolkebeck, 2000a. The effect of various dietary level of phytase and available phosphorus on performance of laying hens. *Poul. sci.*, 79:535-538.
- Boling, S. D., M. W. Douglas, X. Wang, and C. M. Parsons, 2000b. The effect of dietary available phosphorus and phytase on performance of young and old laying hens. *Poul. sci.*, 79:224-230.
- Broz, J., P. Oldale, A. H. Perrin-voltz, G. Tychen, J. Schulze, and C. Simoes Nunes, 1994. Effects of supplemental phytase on performance and phosphorus utilization in broiler chickens fed a low phosphorus diet without addition of inorganic phosphates. *Bri. Poul. sci.*, 35: 273–280.
- Can, A, M. Avci, N. Denek and O. Kaplan, 2007. Effects of microbial phytase on growth performance, carcass yield, biochemical parameters, oxidative stress and fecal phosphorus content of Japanese quails. *J. Anim Vet. Adv.*, 6: 1031-1035.
- Coelho, M.B. and E. Kornegay, 1996. Phytase in Animal Nutrition and Waste Management. BASF Corp., Mount Olive, New Jersey, USA.
- Deutsche Gesellschaft für klinische Chemie, 1972. Empfehlungen der deutschen Gesellschaft für Klinische Chemie (DGKC). Standardisierung von Methoden zur Bestimmung von Enzymaktivitäten in biologischen Flüssigkeiten. (Recommendation of the German Society of Clinical Chemistry. Standardization of methods for measurement of enzymatic activities in biological fluids.) *Z Klin Chem Klin Biochem*, 10:182-92.
- Frost, T. J., and D. A. Roland, Sr, 1991. The influence of various calcium and phosphorus levels on tibia strength and egg shell quality of pullets during peak production. *Poul. sci.*, 70:1640-1643.
- Hall, L. E., R. B. Shirley, R. I. Bakalli, S. E. Aggrey, G. M. Pesti, H. M. Edwards, Jr. 2003. Power of Two Methods for the Estimation of Bone Ash of Broilers. *Poul. sci.*, 82:414–418.
- Huff, W. E., P. A. Moore JR., P. W. Waldroup, A. L. Waldroup, J. M. Balog, G. R. Huff, N. C. Rath, T. C. Daniel and V. Raboy, 1998. Effect of Dietary Phytase and High Available Phosphorus Corn on Broiler Chicken Performance. *Poul. sci.*, 77:1899–1904.

International Union of Biochemistry, 1979. Pp. 242-247 in: Enzyme Nomenclature: Recommendations of the Nomenclature Committee of the International Union of Biochemistry. Academic Press, New York, NY.

Jalal, M. A., and S. E. Scheideler, 2001. Effect of supplementation of two different sources of phytase on egg production parameters in laying hens and nutrient digestibility. *Poul. sci.*, 80:1463-1471.

Jongbloed, A. W., and P. A. Kemme, 1990. Effect of pelleting mixed feeds on phytase activity and the apparent absorbability of phosphorus and calcium in pigs. *Anim. feed Sci. Techn.*, 28:233-242.

Keshavarz, K., 2003. The effect of different levels of nonphytate phosphorus with and without phytase on the performance of four stains of laying hens. *Poul. sci.*, 82:71- 91.

Ketaren, P. P., E. S. Batterham, E. B. Dettmann, and D. J. Farrell, 1993. Phosphorus studies in pigs. 3. Effect of phytase supplementation on the digestibility and availability of phosphorus in soya-bean meal for grower pigs. *Bri. Jou. Nut.*, 70:289-311.

Nation Research Council, 1994. Nutrient Requirements of poultry 9th rev. ed. National Academy Press Washington DC.

Nelson, T. S., T. R. Shieh, R. J. Wodzinski, and J. H. Ware, 1968. The availability of phytate phosphorus in soybean meal before and after treatment with a mold phytase. *Poul. sci.*, 47:1842-1848.

Perney, K. M., A. H. Cantor, M. L. Straw, and K. L. Herkelman, 1993. The effect of dietary phytase on growth performance and phosphorus utilization of broiler chickens. *Poul. sci.*, 72: 2706- 2714.

Qian, H., H. P. Veit, E. T. Kornegay, V. Ravindran, and D. M. Denbow, 1996. Effects of supplemental phytase and phosphorus on histological and other tibia bone characteristics and performances of broilers fed semi-purified diets. *Poul. sci.*, 75: 618-626.

Qian, H., E. T. Kornegay, and D. M. Denbow, 1997. Utilization of phytate phosphorus and calcium as influenced by microbial phytase, cholecalciferol, and the calcium:total phosphorus ratio in broiler diets. *Poul. sci.*, 76: 37-46.

Rama-Rao, S. V., V. Ravindra Reddy, and V. Ramasubba Reddy, 1999. Enhancement of phytate phosphorus availability in the diets of commercial broilers and layers. *Anim. feed Sci. Techn.*, 79:211-222.

Ravindran, V., W. L. Bryden, and E. T. Kornegay, 1995. Phytates: Occurrence bioavailability and implications in poultry nutrition. *Poul. Avi. Biol. Rev.*, 6:125-143.

Ravindran, V., W. L. Bryden, S. Cabahug, and P. H. Selle, 1998. Impact of microbial phytase on the digestibility of protein, amino acids and energy in broilers. Pages 156- 165 in: Proceedings of the Maryland Nutrition Conference for Feed Manufacturing, Baltimore, MD.

Roland, D. A. Sr., H. A. Ahmad, S. S. Yadalam, and T. Sefton, 2003. Effect of nongenetically modified phytase supplementation on commercial Leghorns. *Jou. App. Poul. Res.*, 12:257-263.

Sarcicek, B. Z., Ü. Klc, A. V. Garipoglu, 2005. Replacing soybean meal (SBM) by canola meal (CM): the effects of multi-enzyme and phytase supplementation on the performance of growing and laying quails. *Asian-Australasian Jou. of Anim. Sci.*, www.cababstractsplus.org/abstracts/Abstract.aspx?AcNo=20053172489

SAS Institute, 1998. SAS User's Guide: Statistics. Version 8 edition SAS Institute Inc., Cary, NC.

Sebastian, S., S. P. Touchburn, E. R. Chavez, and P. C. Lague, 1996a. The effects of supplemental microbial phytase on the performance and utilization of dietary calcium, phosphorus, copper, and zinc in broiler chickens fed cornsoybean diets. *Poul. sci.*, 75:729–736.

Sebastian, S., S. P. Touchburn, E. R. Chavez, and P. C. Lague, 1996b. Efficacy of supplemental microbial phytase at different dietary calcium levels on growth performance and mineral utilization of broiler chickens. *Poul. sci.*, 75: 1516–1523.

Sebastian, S., S. P. Touchburn, and E. R. Chavez, 1998. Implication of phytic acid and supplemental microbial phytase in poultry nutrition: A review. *World's Poul. sci. Jou.*, 54:27-47.

Simons, P.C.M., H.A.J. Versteegh, A. W. Jongbloed, P. A. Kemme, P. Slump, K. D. Bos, M.G.E. Wolters, R. F. Beudeker, and G. J. Verschoor, 1990. Improvement of phosphorus availability by microbial phytase in broilers and pigs. *Bri. Jou. Nut.*, 64:525–540.

Summers, J. D., 1995. Reduced dietary phosphorus levels for layers. *Poul. Sci.*, 74:1977- 1983.

Usayran, N., and D. Balnave, 1995. Phosphorus requirements of laying hens fed on wheat based diets. *Br. Poul. sci.*, 36:285-301.

Viveros A., A. Brenes, I. Arija, and C. Centeno, 2002 Effects of Microbial Phytase Supplementation on Mineral Utilization and Serum Enzyme Activities in Broiler Chicks Fed Different Levels of Phosphorus. *Poul. sci.*, 81:1172–1183.

Yi, Z., E.T. Komegay, V. Ravlndran andDM Denbow, 1996. Improving phytate phosphoms availability in com and soybean meal for broiler using microbial phytase and calculation of phosphom equivalency values for phytase *Poult. Sci.*, 75: 240-249.

Zhu, X. S., P. A. Seib, G. L. Allee, and Y. T. Liang, 1990. Preparation of a low-phytate feed mixture and bioavailability of its phosphorus to chicks. *Anim. feed Sci. Techn.*, 27:341–351.